A Linear Mixed-Effects Models approach to obtain Rasch-like estimates of accuracies and response times from fully-crossed design data

Rasch 2.0


Ottavia M. Epifania University of Trento ottavia.epifania@unitn.it


AIP 2025, Torino

September, 12, 2025

The end








Fully-crossed structures

- The "natural" one (so called *compatible* condition)
 - I love Coke and its easier to associate these stimuli to positive attributes
- The "innatural" one (so *incompatible* condition)
 - I love Coke and its harder to associate these stimuli to negative stimuli

Fully-crossed structures

- 1 The "natural" one (so called *compatible* condition)
 - I love Coke and its easier to associate these stimuli to positive attributes
- ② The "innatural" one (so *incompatible* condition)

I love Coke and its harder to associate these stimuli to negative stimuli



The end

Person-level scores

$$s_p = \frac{\bar{X}_{p, \text{comp}} - \bar{X}_{p, \text{inc}}}{s d_{\text{pooled}}}$$

Scoring

Person-level scores

$$s_p = \frac{\bar{X}_{p, \text{comp}} - \bar{X}_{p, \text{ineq}}}{s d_{\text{pooled}}}$$



Advantages

Ease of computation Ease of interpretation Scoring


Person-level scores

$$s_p = \frac{\bar{X}_{p, \mathsf{comp}} - \bar{X}_{p, \mathsf{inc}}}{s d_{\mathsf{pooled}}}$$



Advantages

Ease of computation Ease of interpretation



- Being slow (less accurate) in one condition = being fast (or more accurate) in the opposite one: 0 means absence of bias
- 2 All stimuli have the same impact (fixed effects)

A long tradition

i Respondents are random factors

Sampled from a larger population

Need for acknowledging the sampling variability

Results can be generalized to other respondents belonging to the same population

A long tradition

i Respondents are random factors

Sampled from a larger population Need for acknowledging the sampling variability

Results can be generalized to other respondents belonging to the same popu-

lation

i Stimuli/items are fixed factors

Taken to be entire population

There is no sampling variability

There is no need to generalize the results because the stimuli are the population

- Generalization of the results is impaired
- Error variance everywhere, left free to bias everything
- The information at the stimulus level is lost

- Generalization of the results is impaired
- Error variance everywhere, left free to bias everything
- The information at the stimulus level is lost.

Linear Mixed Effects Models

Rasch model

- Generalization of the results is impaired
- Error variance everywhere, left free to bias everything
- The information at the stimulus level is lost

 \sum

Linear Mixed Effects Models

 ψ

Rasch model



Rasch-like parametrization estimated with Linear Mixed Effects Models

Statistics meets Psychometerics

i Rasch

$$P(x_{ps}=1) = \frac{\exp(\theta_p - b_s)}{1 + \exp(\theta_p - b_s)}$$

$$P(x_{ps}=1) = \frac{\exp(\theta_p \,+\, b_s)}{1 + \exp(\theta_p \,+\, b_s)}$$

Statistics meets Psychomeerics

i Rasch

$$P(x_{ps}=1) = \frac{\exp(\theta_p - b_s)}{1 + \exp(\theta_p - b_s)}$$

GLM (inverse function)

$$P(x_{ps}=1) = \frac{\exp(\theta_p \,+\, b_s)}{1 + \exp(\theta_p \,+\, b_s)} \label{eq:psi}$$

i Log-normal

$$E(t_{ps}) = \delta_s - \tau_p$$

$$E(t_{ps}) = \delta_s + \tau_p + \varepsilon$$

Real data

The end

The proposed workaround

Fully-crossed structures

X: Model Matrix

Needs to be extended:

 β : Coefficients

In a I M:

 $d\!\!:$ Random effects associated to the random factors in Z ... Not model parameters! Best Linear Unbiased Predictors

 $n = \mathbf{X}\beta + \mathbf{Z}d$

 Γ : Parameters estimated for the random factors in the model (variances and covariances)

The end

Random structures

i Models

Model 1

 $y = \beta_c X_c + \alpha_{p[i]} + \alpha_{s[i]}$

$$y = eta_c X_c + lpha_{p[i]} + eta_{s[i]} c_i$$
 Model 3

$$y = \beta_c X_c + \beta_{p[i]} c_i + \alpha_{s[i]}$$



Parametrizations

respondents

GLMM LMM **Model 1**

 $egin{array}{cccc} {
m stimuli} & b_{sc} & \delta_{sc} \ & {
m Model 3} \ \end{array}$

 $\begin{array}{lll} {\rm respondents} & \theta_{pc} & \tau_{pc} \\ {\rm stimuli} & b_s & \delta_s \end{array}$

 $p=1,\ldots,P$: Respondent, $s=1,\ldots,S$: Stimulus, $c\in\{0,1\}$ Associative condition, i Trial

The lower the value, the better the model

! AIC, BIC, and model complexity:

Total number of parameters: β and Γ NOT the levels in d


Model 2 and Model 3: Same complexity, different focus

The chosen model is the least wrong model given the considered models


12 Object stimuli

Fully-crossed structures

White people faces



Black people faces



16 Attribute stimuli

Positive attributes

Good, laughter, pleasure, glory, peace, happy, joy, love

Negative attributes

Evil, bad, horrible, terrible, nasty, pain, failure, hate

Best Fitting Models

GI MMs Model 2 b_{WGBB} and b_{BGWB}

The IAT effect is mostly due to variations in the *stimuli functioning* between conditions, while the performance of the respondents seems unaltered


I MMs Model 3 au_{WGBB} and au_{RGWR}

Real data

The IAT effect is mostly due to variations in the performance of the respondents between conditions, while the functioning of the stimuli appears not affected


Rasch-like estimates

 θ_p




Rasch-like estimates

$b_{\rm WGBB}$ and $b_{\rm WGBB}$




Log-normal estimates


$\tau_{\rm WGBB}$ and $\tau_{\rm BGWB}$



Log-normal estimates

 δ_s





The sky is the limit... but do not over complicate things

- The best model depends on the other models... sometimes useful, never right
- The sky is the limit... but do not over complicate things

HOWEVER

• Time and accuracy are independent from one another, pretty bold assumption

- The best model depends on the other models... sometimes useful, never right
- The sky is the limit... but do not over complicate things

Fully-crossed structures

HOWEVER

Time and accuracy are independent from one another, pretty bold assumption



A Guided Tutorial on Linear Mixed-Effects Models for the Analysis of Accuracies and Response Times in Experiments With Fully Crossed Design

> Ottavia M. Epifania, Pasquale Anselmi, and Egidio Robusto Department of Philosophy, Sociology, Education and Applied Psychology, University of Padova

https://doi.org/10.1037/met0000708